


Central & Northern North Sea Digital Core Log Database



Calderdale Geoscience (CGL), an independent geosciences consultancy established in 2004, has produced a suite of mapping and database products focussed on the Central and Northern North Sea (CNS/NNS).


In preparation for the UK 30th Round, **CGL** can offer CNS/NNS digital core logs. This product facilitates the mapping and modelling of reservoir properties across the UK 30th Round SNS indicative area.



CNS and NNS Cored Intervals:

CGL can offer digital products for the wells indicated (left). **CGL** has evaluated the cored intervals in the following (and other) stratigraphic intervals:

- Tertiary (c. 454 wells)
- Upper Cretaceous (c. 87 wells)
- Lower Cretaceous (c. 186 wells)
- Upper Jurassic (c. 615 wells)
- Middle Jurassic (c. 371 wells)
- Lower Jurassic (c. 140 wells)
- Triassic (c. 230 wells)



Available Products:


CGL core logs can be purchased singly or as groups of wells in the following formats:

- Bespoke drafted images, with wireline, porosity/permeability and petrography data added
- Data-tables of core attributes, both descriptive and interpretative-based, presented as workstation-ready LAS curves and/or spreadsheet interval tables.

SERIES	CHRONO STRATIGRAPHY	GROUP	LITHO STRATIGRAPHY	SEQUENCE STRATIGRAPHY	CORE DEPTH (m)	CORE INTERVAL	SEDIMENTARY STRUCTURES AND GRAIN SIZE		COMMENTS	STANDARD FACIES	SEDIMENTOLOGICAL INTERPRETATION
							GRAPHIC LITHOLOGY	CARBONATE CEMENT			
LATE PALAEOCENE	THANETIAN	MONTROSE	S9	CORE #1	8700	8700-8700	CLAY		CORE 1 GOOD CORE ALL CORE DEPTHS - LOG DEPTH NOT CALIBRATED. MODERATELY SORTED. SANDY SILTY SANDS WITH SANDY SILTY SANDS. SANDY SILTY SANDS WITH SANDY SILTY SANDS.	S _{mc}	SUBMARINE FAN FRINGE: MINOR CHANNEL ATTACHED TURBIDITE LOBE
					8700	8700-8700	SILT		UFF TO YELLOW GREY HOMOGENEOUS FINE TO MEDIUM SANDY SILTY SANDS WITH SANDY SILTY SANDS. SANDY SILTY SANDS WITH SANDY SILTY SANDS.	S _{mb}	INTERLOBE SILTSTONE FAN FILL ABANDONMENT
					8700	8700-8700	FINE SAND		UFF TO OLIVE GREY SILTSTONE WITH DEFORMED VERY FINE SANDY SILTY SANDS.	F _{so}	SUBMARINE FAN FRINGE: MINOR PROGRADING CHANNEL ATTACHED TURBIDITE LOBE
					8700	8700-8700	COARSE SAND		UFF TO YELLOW GREY MODERATELY SORTED SANDSTONE. LARGE LIGHT LENSES IN PLACES.	S _{ld}	GRADUAL LOBE ABANDONMENT
					8700	8700-8700	VERY FINE SAND		UFF TO YELLOW GREY HOMOGENEOUS CARBONACEOUS SANDS TO BUFF CARBONATE CEMENTED SANDSTONE.	F _{ts}	GRADUAL LOBE ABANDONMENT
					8700	8700-8700	COARSE SAND		CLAY GREY SILTSTONE AND VERY FINE GRAINED CARBONACEOUS SANDSTONE.	S _{gn}	SUBMARINE FAN FRINGE: AMALGAMATED CHANNEL FILL TURBIDITES
					8700	8700-8700	FINE SAND		UFF TO YELLOW GREY HOMOGENEOUS SANDSTONE.	F _l	GRADUAL LOBE ABANDONMENT
					8700	8700-8700	COARSE SAND		UFF, HOMOGENEOUS, MODERATELY TO WELL SORTED SANDSTONE.	S _{ld}	SUBMARINE FAN FRINGE: PROGRADING SAND-DOMINATED TURBIDITE LOBE
					8700	8700-8700	VERY FINE SAND		UFF, CARBONACEOUS SANDSTONE.	S _{gn}	
					8700	8700-8700	COARSE SAND		UFF, FINE TO MEDIUM GRAINED SANDSTONE BECOMING MORE CARBONACEOUS TOWARDS TOP OF CORE.	F _{ts}	
					8700	8700-8700	COARSE SAND		CLAY GREY TO GREY SILTSTONE, WITH CONVOLUTED LAMINAE.	F _{ld}	
					8700	8700-8700	COARSE SAND		UFF SANDSTONE WITH LARGE MUDSTONE RIP-UP CLASTS TOWARDS TOP OF SANDSTONE.	S _{mc}	
					8700	8700-8700	COARSE SAND		UFF, WELL SORTED, HOMOGENEOUS SANDSTONE. FINE CHANNEL LAMINAE, TOWARDS BASE.	S _l	
					8700	8700-8700	COARSE SAND		UFF, WELL SORTED, HOMOGENEOUS SANDSTONE.	S _{gn}	
					8700	8700-8700	COARSE SAND		SILTSTONE WITH DEFORMED SANDSTONE LAMINAE.	F _{ld}	

For more information, please contact Paul Emerson; paule@calderdalegeoscience.co.uk
Visit www.calderdalegeoscience.co.uk





Important rock-property data often remains undiscovered and under-utilised in legacy core logs, core reports and well reports (1). **CGL** offers the digital capture of these core attribute data to clients as a bureau service.




The captured digital data can be output as spreadsheets (2) and/or LAS curves (3). These can be imported into workstations for reservoir property analysis and geophysical model calibration. In this example, the original images (1) have been vectorised into descriptive and interpretative curves (2 and 3) as follows:

- Lithofacies code curve with an ordinal scale
- Grain size curve with a Wentworth scale
- Cementation index with an ordinal scale
- Shale (%) curve
- Porosity/permeability point data
- Interpreted facies associations with an ordinal scale


This regularly sampled data can readily be analysed and plotted as distributions (4).

2.

TOP	BOTTOM	LITHOFACIES CODE	PROCESS	LITHOLOGY DOM	STRUCTURE DOM	STRUCTURE SUB	MEDIAN GRAIN SIZE	CLAY CONTENT	CEMENT	COLOUR	
3911.60	3911.80	Ssm	Ff	Ss		b	fU	0	F	LtRd	
3911.80	3911.90	Ssm	Ff	Ss		b	xl	fU	0	F	LtRd
3911.90	3911.95	Ssm	Ff	Ss	C	b	xl	fU	0	F	LtRd
3911.95	3912.00	Ssm	Ff	Ss		b	xl	fL	0	M	LtRd
3912.00	3912.05	Ssm	Ff	Ss		b	xl	mU	0	M	LtRd
3912.05	3912.10	Snf	Ff	Ss	C	pl	ic	fU	0	M	LtRd
3912.10	3912.15	Snf	Ff	Ss	C	pl	ic	fL	0	M	LtRd
3912.15	3912.25	Ssm	Ff	Ss		l	ic	vfU	0	M	LtRd
3912.25	3912.35	Ssm	Ff	Ss		l	xl	vfU	0	M	LtRd
3912.35	3912.40	Ssm	Ff	M	Ss	l	M	60	M	LtRd	
3912.40	3912.50	Ssm	Ff	Ss		l	vfU	0	M	LtRd	
3912.50	3912.55	M	Ff	M		l	M	90	H	DkRd	
3912.55	3912.70	Ssf	Ff	Ss		pl	vfL	0	M	LtRd	
3912.70	3912.80	Ssf	Ff	Ss		b	vfU	0	M	LtRd	
3912.80	3913.00	Ssm	Ff	Ss		b	xh	mL	0	F	LtRd
3913.00	3913.05	Ssf	Ff	Ss		b	xl	fL	0	M	LtRd
3913.05	3913.20	Ssf	Ff	Ss		b	fL	0	M	LtRd	
3913.20	3913.30	Snm	Ff	Ss		h	ic	mU	0	F	LtRd
3913.30	3913.60	Snm	Ff	Ss		h	mL	0	M	LtRd	
3913.60	3914.70	Sw	Aldp	Ss	M	b	w f	fU	20	F	DkRd
3914.70	3914.75	SM	Aldp	M	Ss	l	M	70	H	DkRd	
3914.75	3914.85	Ssf	Ff	Ss		l	vfL	0	F	LtRd	
3914.85	3914.90	Ssf	Ff	Ss		l	xh	fL	0	F	LtRd
3914.90	3914.95	Ssf	Ff	Ss		l	mL	0	M	LtRd	
3914.95	3915.05	SM	Ff	M	Ss	h	sd	M	70	H	DkRd
3915.05	3915.33	Ssf	Ff	Ss		b	xh	fL	0	M	LtRd
3915.33	3915.55	Ssm	Ff	Ss		b	xh	mL	0	F	LtRd
3915.55	3915.75	Ssf	Ff	Ss		b	fL	0	M	LtRd	




For more information, please contact Paul Emerson; paule@calderdalegeoscience.co.uk
 Visit www.calderdalegeoscience.co.uk





In the case of substantial cored intervals, facies associations can be displayed at seismic scale. Here, in an example from the Southern North Sea, an interpreted **facies association curve**, captured from a thick cored section through Rotliegend and Upper Carboniferous strata, has been scaled to emphasise energy of deposition and potential reservoir quality (see below). Excursions of the curve to the left reflect higher energy and better reservoir quality (e.g. channel and mouthbar associations). In addition, curves have been constructed to illustrate the **cored versus un-cored interval** (with a binary 0 (uncored) to 1 (cored) scale) and **depositional environment** (scaled to emphasise more marine environments with excursions to the left). Facies association interpretation can be extrapolated to non cored intervals using wireline data and can also be provided by **CGL**. Interpretation of cored facies association and depositional setting can be used in an accurate prediction of Gross Depositional Environment (GDE) Mapping and Play Fairway Analysis



In the Rotliegend (**BZ**), displaying facies associations at seismic scale, facilitates seismic interpretation:

- Seismically -transparent desert lake/marginal sabkha facies of the Silverpit Clay interval form part of the regional sealing unit.

The Upper Carboniferous (Namurian (**TN**) and Westphalian (**TW**)) section shows a direct and clear relationship between facies, depositional setting, stratigraphy and seismic response:

- Bright reflectors indicating delta top coals and fluvial channel development in upper part of Westphalian unit .
- Development of thick Kinderscoutian ((**TK**) Namurian) deltaic channel sands near base of cored section.
- Gas is encountered in the channelised intervals sourced from adjacent coal-prone delta top.

For more information, please contact Paul Emerson; paule@calderdalegeoscience.co.uk
Visit www.calderdalegeoscience.co.uk

